

TECHNICAL REPORT ARBRL-TR-02059

A COMPUTATIONAL ALGORITHM FOR THE
EIGENVECTORS OF A SINGULAR MATRIX

Palmer R. Schlegel

TECHNICAL
LIBRARY

May 1978



US ARMY ARMAMENT RESEARCH AND DEVELOPMENT COMMAND
BALLISTIC RESEARCH LABORATORY
ABERDEEN PROVING GROUND, MARYLAND

Approved for public release; distribution unlimited.

Destroy this report when it is no longer needed.
Do not return it to the originator.

Secondary distribution of this report by originating
or sponsoring activity is prohibited.

Additional copies of this report may be obtained
from the National Technical Information Service,
U.S. Department of Commerce, Springfield, Virginia
22161.

The findings in this report are not to be construed as
an official Department of the Army position, unless
so designated by other authorized documents.

*The use of trade names or manufacturers' names in this report
does not constitute indorsement of any commercial product.*

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER TECHNICAL REPORT ARBRL-TR-02059	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A COMPUTATIONAL ALGORITHM FOR THE EIGENVECTORS OF A SINGULAR MATRIX.		5. TYPE OF REPORT & PERIOD COVERED Final
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Palmer R. Schlegel		8. CONTRACT OR GRANT NUMBER(s)
9. PERFORMING ORGANIZATION NAME AND ADDRESS U.S. Army Ballistic Research Laboratory (ATTN: DRDAR-BLB) Aberdeen Proving Ground, MD 21005		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS RDTE Project 1L161101A91A
11. CONTROLLING OFFICE NAME AND ADDRESS U.S. Army Armament Research and Development Command U.S. Army Ballistic Research Laboratory (ATTN: DRDAR) Aberdeen Proving Ground, MD 21005		12. REPORT DATE MAY 1978
		13. NUMBER OF PAGES 19
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) eigenvalue, eigenvector, singular matrix, nonsingular matrix, computational algorithm		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In this paper a computational, non-iterative type algorithm is given to obtain the eigenvectors of a singular matrix. The algorithm yields a (smaller) nonsingular matrix that has exactly the same nonzero eigenvalues as the given singular matrix and only these eigenvalues. The eigenvectors for the singular matrix are constructed from the eigenvectors of the nonsingular matrix. The eigenvectors associated with the zero eigenvalues of the singular matrix are an immediate consequence of the algorithm.		

TABLE OF CONTENTS

I.	INTRODUCTION.	5
II.	MOTIVATION OF THE ALGORITHM	5
III.	EXISTENCE OF THE MATRIX U	7
IV.	CONSTRUCTION OF THE MATRIX U.	9
V.	SOME ILLUSTRATIVE EXAMPLES.	11
VI.	CONCLUSION.	16
	ACKNOWLEDGEMENT	17
	DISTRIBUTION LIST	19

I. INTRODUCTION

In this paper the eigenvalues and the eigenvectors of a singular matrix, A , are investigated.¹ Present techniques for determining the eigenvalues of a matrix require that the matrix be nonsingular or that some iterative procedure such as root searching techniques be applied to the characteristic equation. Here, a computational (non-iterative type) algorithm is given to construct a (smaller) nonsingular matrix, Q , that has exactly the same nonzero eigenvalues as A and only these eigenvalues. Thus, a standard technique such as the QR algorithm² can be applied to the matrix Q to evaluate the eigenvalues and eigenvectors of Q .

The algorithm given in this paper yields the eigenvectors of the nonzero eigenvalues of A , which are obtained from the eigenvectors of Q , and the rank of the matrix A . The eigenvectors associated with the zero eigenvalues of A are an immediate consequence of the algorithm.

II. MOTIVATION OF THE ALGORITHM

Let $A = (a_{ij})$ be an $n \times n$ matrix and U an $n \times n$ nonsingular matrix. Define $W = U^{-1}AU$. Let λ and X be an eigenvalue and an eigenvector, respectively, of A , then

$$\begin{aligned} AX &= \lambda X \\ U^{-1}A(UU^{-1})X &= \lambda U^{-1}X \\ W(U^{-1}X) &= \lambda(U^{-1}X), \end{aligned} \tag{1}$$

that is, λ and $U^{-1}X$ are an eigenvalue and eigenvector of W . Conversely, if μ and Z are an eigenvalue and an eigenvector of W , then μ and UZ are an eigenvalue and an eigenvector of A .

Suppose A is of rank r and there exists a U such that W can be partitioned into the following form:

¹The algorithm given in this paper was developed by the author in response to a problem posed by H. McCoy, TRASANA, to obtain the eigenvalues and eigenvectors of a special singular matrix.

²Wilkinson, J. H., The Algebraic Eigenvalue Problem, Clarendon Press, Oxford, 1965.

$$W = U^{-1}AU = \begin{pmatrix} Q_r & O_1 \\ B & O_2 \end{pmatrix}, \quad (2)$$

where Q_r is an $r \times r$ matrix, O_1 and O_2 are $r \times (n-r)$ and $(n-r) \times (n-r)$ zero matrices, respectively, and B is an $(n-r) \times r$ matrix. Let $\lambda \neq 0$ be an eigenvalue of W , and of A , and Z be the associated eigenvector. If the vector Z , as a matrix, is partitioned into Z_r , a $r \times 1$ matrix, and Y , a $(n-r) \times 1$ matrix, then we can write the following matrix equation

$$\begin{pmatrix} Q_r & O_1 \\ B & O_2 \end{pmatrix} \begin{pmatrix} Z_r \\ Y \end{pmatrix} = \lambda \begin{pmatrix} Z_r \\ Y \end{pmatrix} \quad (3)$$

as

$$Q_r Z_r = \lambda Z_r \quad (4)$$

and

$$BZ_r = \lambda Y, \quad (5)$$

that is, λ and Z_r are an eigenvalue and eigenvector of the matrix Q_r .

Thus, the nonzero eigenvalues of W , therefore for A , are the eigenvalues (nonzero) of Q_r . The associated eigenvectors of A are given by

$$X = UZ = U \begin{pmatrix} Z_r \\ Y \end{pmatrix} = U \begin{pmatrix} Z_r \\ \lambda^{-1} BZ_r \end{pmatrix}, \quad (6)$$

which are determined from the eigenvectors of Q_r .

Conversely, if $\lambda \neq 0$ is an eigenvalue of Q_r and Z_r the eigenvector, then it follows that X , which is defined by (6), is an eigenvector of A and λ is the eigenvalue.

The eigenvectors of A associated with the zero eigenvalues are immediate. Define the column vectors

$$E_i = \begin{pmatrix} \delta_{1,r+i} \\ \delta_{2,r+i} \\ \vdots \\ \delta_{n,r+i} \end{pmatrix}, \quad i = 1, 2, \dots, n-r, \quad (7)$$

where δ_{kl} is the Kronecker delta function. Then $WE_i = 0$, that is, E_i , $i = 1, 2, \dots, n-r$, are the associated eigenvectors for the zero eigenvalues of W . Thus, UE_i , $i = 1, 2, \dots, n-r$, are the eigenvectors of the zero eigenvalues of A ; but these eigenvectors are just the column vectors represented by the last $n-r$ columns of the matrix U .

Since the rank of A is r , it follows that $\{UE_i\}$ is the total set of (distinct) eigenvectors associated with the zero eigenvalue. If Q_r is singular, this implies that the characteristic equation has a zero of multiplicity greater than $n-r$; but no greater collection of eigenvectors. The process would then be applied to Q_r to determine a matrix of smaller order that has the same nonzero eigenvalues as A .

III. EXISTENCE OF THE MATRIX U

We will now show the existence of the matrix U . Let A_i , $i = 1, 2, \dots, n$, denote the column vectors represented by the columns of the matrix A . The definition and notation for the inner product of any two vectors is given by

$$(A_i, A_j) = \sum_{k=1}^n a_{ki} \bar{a}_{kj}, \quad (8)$$

where \bar{a} denotes the complex conjugate of a .

We will now apply the Gram-Schmidt orthogonalization process (see [3]) to the set of vectors $\{A_i\}$. If A_1 is the zero vector, then interchange A_1 and A_n . Let U_1 be the elementary matrix that interchanges column one and column n , when post matrix multiplication is applied (this is the identity matrix with the first and n th columns interchanged). If, after this interchange, A_1 is zero, then interchange A_1 and A_{n-1} , where U_2 is the appropriate elementary matrix. Continue until A_1 is not the zero vector. Define $V_1 = A_1$ and

³ Berberian, S. K., Introduction to Hilbert Space, Oxford University Press, New York, 1961.

$$V_2 = A_2 - \alpha_{21}V_1, \quad (9)$$

where $\alpha_{21} = \frac{(A_2, V_1)}{(V_1, V_1)}.$

If V_2 is the zero vector, interchange A_2 and A_k , where A_{k+1} was the last vector interchanged; and U_j the appropriate elementary matrix. Define U_{j+1} to be the elementary matrix that adds $-\alpha_{21}$ times the second column to column k . This is given by appending to the identity matrix $-\alpha_{21}$ in row 2 and column k . Construct a new V_2 . If V_2 is not the zero vector, then continue in the construction until a zero vector is generated, that is, define

$$V_i = A_i - \sum_{j=1}^{i-1} \alpha_{ij} V_j, \quad (10)$$

where $\alpha_{ij} = \frac{(A_i, V_j)}{(V_j, V_j)}, j = 1, \dots, i-1$. If V_i is the zero vector, interchange A_i with the last vector not interchanged, say A_m , and assign an appropriate elementary matrix U_k . Since $V_j, j = 1, \dots, i-1$, can be written in terms of A_1, \dots, A_j , then A_i can be written in the following form:

$$A_i = \beta_{i1}A_1 + \dots + \beta_{i,i-1}A_{i-1}. \quad (11)$$

Furthermore, let $U_{k+j}, j = 1, \dots, i-1$, be the elementary matrices which adds $-\beta_{ij}$ times column j to column m . Continue this construction for V_i until A_{i+1} was the last vector interchanged.

Define U to be the product of the elementary matrices generated above, that is,

$$U = U_1 U_2 \dots U_t. \quad (12)$$

Then AU has the form:

$$AU = \begin{pmatrix} Q & 0_1 \\ R & 0_2 \end{pmatrix}. \quad (13)$$

For example, if V_i is the zero vector, then

$$A_i = \sum_{j=1}^{i-1} \beta_{ij} A_j . \quad (14)$$

The product of the associated elementary matrices, say U_k, \dots, U_{k+i-1} , would interchange column i with column m and subtract β_{ij} times column j , $j = 1, \dots, i-1$, from column m . This would result in a zero column in column m .

If U_p interchanges column i with column j , then premultiplication by U_p^{-1} would interchange row i with row j ; and if U_q would add $-\beta$ times column i to column j , U_q^{-1} would add β times row j to row i . Thus,

$$U^{-1} = U_t^{-1} \dots U_2^{-1} U_1^{-1} \quad (15)$$

would result in row operations. Therefore, $U^{-1}AU$ would have the same form as (13), that is,

$$U^{-1}AU = \begin{pmatrix} Q_r & 0_1 \\ B & 0_2 \end{pmatrix} . \quad (16)$$

The resulting collection of nonzero vectors, A_1, \dots, A_r , under the Gram-Schmidt process, is the largest collection of independent vectors represented by the columns of A ; therefore, r is the rank of A .

IV. CONSTRUCTION OF THE MATRIX U

In this section we will develop a computationally feasible algorithm for the construction of the matrices U , Q_r and B . It should be noted that the actual construction of U^{-1} will not be required to obtain the final result.

To start the construction, set $U = (u_{ij}) = (\delta_{ij})$, the identity matrix. If there exists a vector V_i equal to the zero vector, then from (14)

$$A_i = \sum_{j=1}^{i-1} \beta_{ij} A_j . \quad (18)$$

Assume A_{m+1} was the last column interchanged. If $u_{ii} = 1$, set $u_{ii} = 0$, $u_{im} = 1$, $u_{mi} = 1$ and $u_{mm} = 0$. If $u_{ii} = 0$, set $u_{m+1,i} = 0$, $u_{m+1,m} = 1$, $u_{mi} = 1$ and $u_{mm} = 0$. This interchanges column i and column m .

Simultaneously, replace column i with column m in matrix A . Note that column m need not be replaced by column i , for this column is assumed to be the zero vector. In order to accomplish this, add $-\beta_{ij}$,

$j = 1, \dots, i-1$, to column m in the U matrix, where the i th row is determined from the one and only one nonzero element in the j th column of the U matrix. This nonzero element is unity. This construction is continued for i until $i+1$ was the last column interchanged. Thus, A has been reduced to the form of (13).

In order to obtain Q_r and B , similar operations must be done on rows of the reduced matrix A . An accounting must be kept on the column operations, for the row operations must be done in the same order, that is, if A has been postmultiplied by $U_1 U_2 \dots U_t$, then A must be pre-multiplied by $U_t^{-1} \dots U_2^{-1} U_1^{-1}$. Note that the row operations only involve the first r columns, for the remaining $n-r$ columns are assumed to be zero vectors. Therefore, if a column operation involved interchanging column ℓ with column m , then the row operation would interchange row ℓ with row m . Similarly, if a column operation adds $-\beta$ times column ℓ to column m , then the row operation would add β times row m to row ℓ .

In order to construct β_{ij} , the vectors, V_i , must be generated, from which β_{ij} can be obtained recursively. From (10)

$$V_i = A_i - \sum_{k=1}^{i-1} \alpha_{ik} V_k , \quad (19)$$

where $\alpha_{ik} = \frac{(A_i, V_k)}{(V_k, V_k)}$. Suppose

$$V_k = A_k - \sum_{j=1}^{k-1} \beta_{kj} A_j , \quad k = 2, \dots, i-1, \quad (20)$$

where $V_1 = A_1$. Then from (19) and (20)

$$\begin{aligned}
V_i &= A_i - \alpha_{i1}A_1 - \sum_{k=2}^{i-1} \alpha_{ik}(A_k - \sum_{j=1}^{k-1} \beta_{kj}A_j) \\
&= A_i - \alpha_{i1}A_1 - \sum_{k=2}^{i-1} \alpha_{ik}A_k + \sum_{k=2}^{i-1} \sum_{j=1}^{k-1} \alpha_{ik}\beta_{kj}A_j \\
&= A_i - \sum_{j=1}^{i-1} \alpha_{ij}A_j + \sum_{j=1}^{i-2} \left(\sum_{k=j+1}^{i-1} \alpha_{ik}\beta_{kj} \right) A_j \\
&= A_i - \alpha_{i,i-1}A_{i-1} - \sum_{j=1}^{i-2} \left(\alpha_{ij} - \sum_{k=j+1}^{i-1} \alpha_{ik}\beta_{kj} \right) A_j. \quad (21)
\end{aligned}$$

Therefore,

$$\beta_{ij} = \begin{cases} \alpha_{ij} - \sum_{k=j+1}^{i-2} \alpha_{ik}\beta_{kj} & , j = 1, \dots, i-2 \\ \alpha_{ij} & , j = i-1 \end{cases} \quad (22)$$

V. SOME ILLUSTRATIVE EXAMPLES

A. Example 1.

Let

$$A = \begin{bmatrix} 1 & 2 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 2 & 3 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

Then

$$V_1 = \begin{bmatrix} 1 \\ 1 \\ 2 \\ 0 \end{bmatrix},$$

which is not the zero vector.

$$V_2 = A_2 - \alpha_{21}V_1 = \begin{bmatrix} 2 \\ 1 \\ 3 \\ 0 \end{bmatrix} - \frac{3}{2} \begin{bmatrix} 1 \\ 1 \\ 2 \\ 0 \end{bmatrix} = \begin{bmatrix} .5 \\ -.5 \\ 0 \\ 0 \end{bmatrix},$$

where

$$\alpha_{21} = \frac{(A_2, V_1)}{(V_1, V_1)} = \frac{2 + 1 + 6 + 0}{1 + 1 + 4 + 0} = \frac{3}{2}$$

and $\beta_{21} = \alpha_{21}$

$$V_3 = A_3 - \alpha_{31}V_1 - \alpha_{32}V_2 = \begin{bmatrix} 0 \\ 1 \\ 1 \\ 0 \end{bmatrix} - \frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ 2 \\ 0 \end{bmatrix} - (-1) \begin{bmatrix} .5 \\ -.5 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix},$$

where

$$\alpha_{31} = \frac{(A_3, V_1)}{(V_1, V_1)} = \frac{0 + 1 + 2 + 0}{6} = \frac{1}{2}$$

$$\alpha_{32} = \frac{(A_3, V_2)}{(V_2, V_2)} = \frac{0 + 0 + (-.5) + 0}{.25 + .25 + 0 + 0} = -1$$

and

$$\beta_{31} = \alpha_{31} - \alpha_{32}\beta_{21} = 2$$

$$\beta_{32} = \alpha_{32} = -1.$$

Now V_3 is the zero vector. Therefore, replace column 3 with column 4 in A . We will assume column 4 is the zero column. For in theory, $-\beta_{31}$ times column 1 and $-\beta_{32}$ times column 2 are added to column 4. Hence, the reduced A matrix is

$$AU = \begin{bmatrix} 1 & 2 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 2 & 3 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

If the same column operations are applied to U, which was initially set to the identity matrix, then

$$U = \begin{bmatrix} 1 & 0 & 0 & -2 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}.$$

Since V_3 was the zero vector, A_3 is renamed (interchange columns). A new V_3 is generated, namely,

$$V_3 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} - 0 \cdot V_1 - 0 \cdot V_2 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}$$

This "new" V_3 is not the zero vector. Hence, column operations terminate. The row operations are the following:

1. Interchange row 3 and row 4.
2. β_{31} times row 4 added to row 1.
3. β_{32} times row 4 added to row 2.

Thus,

$$U^{-1}AU = \begin{bmatrix} 5 & 8 & 0 & 0 \\ -1 & -2 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 2 & 3 & 0 & 0 \end{bmatrix}$$

The rank of A is 3. The eigenvector associated with the zero eigenvalue of A is the fourth column of U. Since

$$Q_r = \begin{bmatrix} 5 & 8 & 0 \\ -1 & -2 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

is nonsingular, no further reduction need be applied. The eigenvalues of Q_r are $\lambda_1 = 1$, $\lambda_2 = \frac{1}{2}(3 + \sqrt{17})$ and $\lambda_3 = \frac{1}{2}(3 - \sqrt{17})$. The respective eigenvectors are the following:

$$Z_1 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, \quad Z_2 = \frac{1}{2} \begin{bmatrix} -7-\sqrt{17} \\ 2 \\ 0 \end{bmatrix}, \quad Z_3 = \frac{1}{2} \begin{bmatrix} -7+\sqrt{17} \\ 2 \\ 0 \end{bmatrix}.$$

To obtain the eigenvectors for the nonzero eigenvalues of A, (6) is applied, where

$$B = \begin{bmatrix} 2 & 3 & 0 \end{bmatrix}.$$

Thus,

$$X_1 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \quad X_2 = \frac{1}{4} \begin{bmatrix} -4 \\ -1-\sqrt{17} \\ -5-\sqrt{17} \\ 0 \end{bmatrix}, \quad X_3 = \frac{1}{4} \begin{bmatrix} -4 \\ -1+\sqrt{17} \\ -5+\sqrt{17} \\ 0 \end{bmatrix}$$

and the eigenvector for the zero eigenvalue is

$$X_4 = \begin{bmatrix} -2 \\ 1 \\ 1 \\ 0 \end{bmatrix},$$

the last column of U.

B. Example 2.

Let

$$A = \begin{bmatrix} 3 & -2 & -1 & 3 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

It is easy to see that A_1 , A_2 and A_3 are independent vectors (also V_1 , V_2 and V_3) and $A_4 = A_1$. Thus, the only column operation is to add -1 times column 1 to column 4. Therefore,

$$Q_T = \begin{bmatrix} 3 & -2 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$

and

$$B = [0 \ 0 \ 1],$$

where

$$U = \begin{bmatrix} 1 & 0 & 0 & -1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

If we apply the algorithm again to Q'_r , without intermediate calculations, we have

$$Q'_r = \begin{bmatrix} 3 & -2 \\ 1 & 0 \end{bmatrix},$$

$$B' = [0 \ 1]$$

and U' is the 3×3 identity matrix, where the prime denotes the partitioning of Q_r . The two eigenvalues of Q'_r are $\lambda_1 = 2$ and $\lambda_2 = 1$, and the associated eigenvectors are

$$z'_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

and

$$z'_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}.$$

The respective eigenvectors of Q_r are

$$z_1 = U' \begin{bmatrix} z'_1 \\ \lambda_1^{-1} B' z'_1 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 4 \\ 2 \\ 1 \end{bmatrix}$$

and

$$z_2 = U' \begin{bmatrix} z'_2 \\ \lambda_2^{-1} B' z'_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}.$$

Finally, the eigenvectors of the nonzero eigenvalues of A are

$$x_1 = U \begin{bmatrix} z_1 \\ \lambda_1^{-1} B z_1 \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 7 \\ 4 \\ 2 \\ 1 \end{bmatrix}$$

and

$$x_2 = U \begin{bmatrix} z_2 \\ \lambda_2^{-1} B z_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 1 \\ 1 \end{bmatrix}.$$

The eigenvector associated with the zero eigenvalue is

$$x_3 = \begin{bmatrix} -1 \\ 0 \\ 0 \\ 1 \end{bmatrix},$$

that is, the fourth column of U . Since the rank of A is three, this is the only eigenvector associated with the zero eigenvalue.

VI. CONCLUSION

In order to apply this algorithm to a computer code, an a priori decision must be made for a "zero vector", due to machine round-off. In the author's program, which generates the matrices U , A_r and B , a vector, V , is the zero vector if

$$(V, V) < \epsilon,$$

where ϵ is some preassigned value.

An alternate application of the procedure could eliminate some computer code "bookkeeping". Since matrix multiplication is associative, a column operation, U_p , could be immediately followed by the row operation U_p^{-1} . This would, of course, increase the arithmetical operations, since the knowledge of the number of zero columns could not be totally incorporated.

ACKNOWLEDGEMENT

The author gratefully acknowledges R. E. Shear, who found some technical errors, and Dr. M. S. Taylor for his many useful editorial comments which were incorporated in this paper.

DISTRIBUTION LIST

<u>No. of</u> <u>Copies</u>	<u>Organization</u>	<u>No. of</u> <u>Copies</u>	<u>Organization</u>
12	Commander Defense Documentation Center ATTN: DDC-TCA Cameron Station Alexandria, VA 22314	1	Commander US Army Mobility Equipment Research & Development Cmd ATTN: DRDME-WC Fort Belvoir, VA 22060
1	Commander US Army Materiel Development and Readiness Command ATTN: DRCDMA-ST 5001 Eisenhower Avenue Alexandria, VA 22333	1	Commander US Army Armament Materiel Readiness Command ATTN: DRSAR-LEP-L, Tech Lib Rock Island, IL 61299
1	Commander US Army Aviation Research and Development Command ATTN: DRSAR-E 12th and Spruce Streets St. Louis, MO 63166	2	Commander US Army Armament Research and Development Command ATTN: DRDAR-TSS (2 cys) Dover, NJ 07801
1	Director US Army Air Mobility Research and Development Laboratory Ames Research Center Moffett Field, CA 94035	1	Commander US Army Harry Diamond Labs ATTN: DRXDO-TI 2800 Powder Mill Road Adelphi, MD 20783
1	Commander US Army Electronics Command ATTN: DRSEL-RD Fort Monmouth, NJ 07703	1	Director US Army TRADOC Systems Analysis Activity ATTN: ATAA-SL, Tech Lib White Sands Missile Range NM 88002
1	Commander US Army Missile Research and Development Command ATTN: DRDMI-R Redstone Arsenal, AL 35809		<u>Aberdeen Proving Ground</u> Marine Corps Ln Ofc Dir, USAMSAA ATTN: Dr. J. Sperrazza Dr. B. Cummings Mr. W. Webster
1	Commander US Army Tank Automotive Research & Development Cmd ATTN: DRDTA-RWL Warren, MI 48090		